



Research Paper

Copyright © Gph rights are reserved by Souad I. Mugassabi

« The Addition and Subtraction Operations for two Continued Fractions »

Souad I. Mugassabi and Fatima F. Abdullah

University of Benghazi, Benghazi, Libya

| | |
|--|------------------------------|
| Corresponding author: * Souad I. Mugassabi | Received Date: July 16, 2020 |
| Tel.: ***** Email: hidden | Published Date: Aug 03, 2020 |

Abstract:

The continued fractions (C.F.) with positive non integer numerators are considered. The addition and subtraction operations of two continued fractions are discovered. Many definitions and examples that we used of these low are presented.

Keywords: Continued fractions, Addition operation of continued fractions, Subtraction operation of continued fractions.

1. INTRODUCTION

Continued fractions have been extensively studied and there is a large body of research related to them [3]. They are now included as classical results in most undergraduate textbooks like [1, 4, 5] and they continue to be studied a great deal. Continued fraction expansions can also be finite or infinite. For example, any rational number can be expressed as a finite continued fraction expansion, while an irrational number has an infinite continued fraction expansion [2, 9]. Although our main concentration is in the operations of the continued fractions, it is important to look for similarities. Previous studies have shown that the operations of the simple continued fractions follow similar patterns to those of continued fractions under certain circumstances [6, 7, and 8]. Therefore it is important to make this comparison. Hence, we summarize some important results for simple continued fractions. The main justification for the paper is the operations of the continued fractions. The addition and subtraction of two continued fractions.

II. PRELIMINARIES

Definition 2.1: A continued fraction is an expression of the form $a_0 + \frac{b_0}{a_1 + \frac{b_1}{a_2 + \frac{b_2}{\ddots + \frac{b_n}{a_n}}}}$ where a_0, a_1, a_2, \dots and b_0, b_1, b_2, \dots can be real or

complex, and their numbers can be either finite or infinite. In this paper, we consider the continued fractions of the form $a_0 + \frac{z}{a_1 + \frac{z}{a_2 + \frac{z}{\ddots + \frac{z}{a_n}}}}$, where a_0 is an integer, $a_1, a_2, a_3, \dots, a_n$ are positive integer and z is a positive non integer. We denote it by $[a_0; a_1, a_2, \dots]_z$.

Example 2.1

$$(a) 3 + \frac{\frac{3}{2}}{4 + \frac{2}{7}} = [3; 4, 7]_{\frac{3}{2}} \quad (b) -7 + \frac{\frac{2}{3}}{1 + \frac{3}{3}} = [-7; 1, 3]_{\frac{2}{3}} \quad (c) 1 + \frac{\frac{5}{3}}{2 + \frac{3}{3 + \frac{5}{3}}} = [1; 2, 3, \dots]_{\frac{5}{3}}$$

(a), (b) have a finite number of terms, and (c) has an infinite number of terms.

Theorem 2.1: A number is rational if and only if it can be expressed as a finite C.F. [8].



Example 2.2:

$$\begin{aligned}
 \text{(a)} \quad \frac{40}{11} &= 3 + \frac{\frac{3}{33}}{\frac{14}{14}} = 3 + \frac{\frac{3}{21}}{\frac{2+\frac{2}{21}}{5}} = 3 + \frac{\frac{3}{15}}{\frac{2+\frac{2}{15}}{\frac{4+\frac{2}{15}}{2}}} = 3 + \frac{\frac{3}{9}}{\frac{2+\frac{2}{9}}{\frac{4+\frac{2}{9}}{\frac{7+\frac{2}{9}}{3}}}} = [3,2,4,7,3]_{3/2}. \\
 \text{(b)} \quad \frac{28}{11} &= 2 + \frac{\frac{2}{11}}{\frac{9}{9}} = 2 + \frac{\frac{2}{6}}{\frac{1+\frac{2}{6}}{3}} = [2,1,3]_{2/3}. \\
 \text{(c)} \quad \frac{3}{7} &= 0 + \frac{\frac{3}{7}}{\frac{2}{2}} = 0 + \frac{\frac{3}{4}}{\frac{3+\frac{3}{4}}{2}} = [0,3,3]_{3/2}. \\
 \text{(d)} \quad \frac{7}{9} &= 0 + \frac{7}{9} = [0;1]_{7/9}.
 \end{aligned}$$

Remark 2.1: To expand a negative rational number $-\frac{a}{b}$ ($a, b > 0$) into C.F we take the greatest integer number $\left\lfloor \frac{-a}{b} \right\rfloor$ for the first term of C.F that is, $\left\lfloor \frac{-a}{b} \right\rfloor = a'_0$, where $-a'_0 \leq -\frac{a}{b} < -a'_0 + 1$. We write $-\frac{a}{b} = -a'_0 + \frac{z}{b'}$ and then we use the same techniques in theorem 2.1 to get the remaining terms for $\frac{z}{b'}$. That is, if $\frac{z}{b'} = [a'_1, a'_2, \dots, a'_n]_z$. Then $-\frac{a}{b} = [a'_0; a'_1, a'_2, \dots, a'_n]_z$. [6].

Example 2.3: $-\frac{71}{11} = -7 + \frac{6}{11} = -7 + \frac{\frac{2}{3}}{\frac{1+\frac{2}{3}}{3}} = [-7,1,3]_{2/3}$.

Lemma 2.1: $[c_0; c_1, \dots, c_{j-1}, 0, c_{j+1}, \dots, c_n]_z = [c_0; c_1, \dots, c_{j-2}, c_{j-1} + c_{j+1}, c_{j+2}, \dots, c_n]_z$

III. ADDITION OPERATION FOR TWO CONTINUED FRACTIONS

Definition 3.1: The C.F $[a_0; a_1, a_2, \dots, a_n]_z$ can be defined by $[a_0; a_1, a_2, \dots, a_n]_z = a_0 + \frac{z k_{n-1}(a_2)}{k_n(a_1)}$ or $[a_0; a_1, a_2, \dots, a_n]_z = \frac{k_{n+1}(a_0)}{k_n(a_1)}$, where $k_i(a_j) = a_{i+(j-1)}k_{i-1}(a_j) + zk_{i-2}(a_j)$, $i = 1, 2, \dots, n, j = 0, 1, \dots, n$ and $k_{-i}(a_j) = 0, k_0(a_j) = 1$.

Definition 3.2: Let $[a_0; a_1, \dots, a_m]_z$ and $[b_0; b_1, \dots, b_n]_z$ be two C.F. we define addition by

(1) If $m = n$ then $[a_0; a_1, \dots, a_n]_z + [b_0; b_1, \dots, b_n]_z = [c_0; c_1, \dots, c_n]_z$ (3.2a)

where

$c_0 = a_0 + b_0 = k_1(a_0) + k_1(b_0)$.

$c_1 = \left\lfloor \frac{a_1 b_1}{a_1 + b_1} \right\rfloor = \left\lfloor \frac{k_1(a_1)k_1(b_1)}{k_1(a_1) + k_1(b_1)} \right\rfloor$.

In general

$$c_m = \begin{cases} \left\lfloor \frac{z(k_{m-1}(a_2)k_m(b_1)k_{m-3}(c_2) - (k_{m-1}(a_2)k_m(b_1) + k_{m-1}(b_2)k_m(a_1))k_{m-2}(c_1))}{(k_{m-1}(a_2)k_m(b_1) + k_{m-1}(b_2)k_m(a_1))k_{m-1}(c_1) - k_m(a_1)k_m(b_1)k_{m-2}(c_2)} \right\rfloor, & m \text{ odd} \\ \left\lfloor \frac{z((k_{m-1}(a_2)k_m(b_1) + k_{m-1}(b_2)k_m(a_1))k_{m-2}(c_1) - k_m(a_1)k_m(b_1)k_{m-3}(c_2))}{k_m(a_1)k_m(b_1)k_{m-2}(c_2) - (k_{m-1}(a_2)k_m(b_1) + k_{m-1}(b_2)k_m(a_1))k_{m-1}(c_1)} \right\rfloor, & m \text{ even} \end{cases}$$

for $m = 2, \dots, n$. The last term c_n of the resulting C.F. is to be expanded again as a C.F.

(2) If $m \neq n$ (suppose that $m < n$) then

$[a_0; a_1, \dots, a_m]_z + [b_0; b_1, \dots, b_m, b_{m+1}, \dots, b_n]_z = [c'_0; c'_1, \dots, c'_m, c'_{m+1}, \dots, c'_n]_z$ (3.2b),

where $c'_j = c_j$ for $j = 0, 1, 2, \dots, m$ (c_j as we did for case $m = n$), while $c'_j = c_{j, j-m}$ and

$$c_{j, j-m} = \begin{cases} \left\lfloor \frac{z(k_{m-1}(a_2)k_j(b_1)k_{j-3}(c'_2) - (k_{m-1}(a_2)k_j(b_1) + k_{j-1}(b_2)k_m(a_1))k_{j-2}(c'_1))}{(k_{m-1}(a_2)k_j(b_1) + k_{j-1}(b_2)k_m(a_1))k_{j-1}(c'_1) - k_m(a_1)k_j(b_1)k_{j-2}(c'_2)} \right\rfloor; & j \text{ odd} \\ \left\lfloor \frac{z((k_{m-1}(a_2)k_j(b_1) + k_{j-1}(b_2)k_m(a_1))k_{j-2}(c'_1) - k_m(a_1)k_j(b_1)k_{j-3}(c'_2))}{k_m(a_1)k_j(b_1)k_{j-2}(c'_2) - (k_{m-1}(a_2)k_j(b_1) + k_{j-1}(b_2)k_m(a_1))k_{j-1}(c'_1)} \right\rfloor; & j \text{ even} \end{cases}$$

for $j = m + 1, m + 2, \dots, n$. Also, the last term of the resulting C.F is to be treated as mentioned earlier.

Example 3.1: Find $[2; 1, 3]_{2/3} + [3; 1, 14]_{2/3}$

Solution: Let $[2; 1, 3]_{2/3} = [a_0; a_1, a_2]_z$ and $[3; 1, 14]_{2/3} = [b_0; b_1, b_2]_z$. We get $m = n = 2$. From definition (3.2a) we get $[a_0; a_1, a_2]_z + [b_0; b_1, b_2]_z = [c_0; c_1, c_2]_z$, where c_2 is the last term and

$c_0 = a_0 + b_0 = 5$.

$c_1 = \left\lfloor \frac{a_1 b_1}{a_1 + b_1} \right\rfloor = \left\lfloor \frac{1}{2} \right\rfloor = 0$.

$c_2 = \frac{z(a_2(b_1 b_2 + z) + b_2(a_1 a_2 + z))}{(a_1 a_2 + z)(b_1 b_2 + z) - (a_2(b_1 b_2 + z) + b_2(a_1 a_2 + z))c_1} = \frac{13}{11}$ (to be treated as C.F),

$$= [1;3,1]_{2/3}$$

$$\text{Then } [2;1,3]_{2/3} + [3;1,14]_{2/3} = [5;0,1,3,1]_{2/3}$$

Example 3.2: Find $[2;2]_{7/4} + [2;9,5,7]_{7/4}$.

Solution: Let $[2;2]_{7/4} = [a_0; a_1]_{7/4}$ and $[2;9,5,7]_{7/4} = [b_0; b_1, b_2, b_3]_{7/4}$, we get $m = 1, n = 3$ and $m < n$, from definition (3.2b) we have $[2;2]_{7/4} + [2;9,5,7]_{7/4} = [a_0; a_1]_{7/4} + [b_0; b_1, b_2, b_3]_{7/4} = [c'_0; c'_1, c'_2, c'_3]_z$, where c'_3 is last term and

$$c'_0 = c_0 = a_0 + b_0 = 2 + 2 = 4.$$

$$c'_1 = c_1 = \left| \frac{a_1 b_1}{a_1 + b_1} \right| = \left| \frac{2 \cdot 9}{2 + 9} \right| = \left| \frac{18}{11} \right| = 1.$$

$$c'_2 = c_{2,1} = \left| \frac{z((k_0(a_2)k_2(b_1) + k_1(b_2)k_1(a_1))k_0(c'_1) - k_1(a_1)k_2(b_1)k_{-1}(c'_2))}{k_1(a_1)k_2(b_1)k_0(c'_2) - (k_0(a_2)k_2(b_1) + k_1(b_2)k_1(a_1)k_1(c'_1)))} \right|$$

$$= \left| \frac{z(((1)(b_1 b_2 + z) + b_2 a_1)(1) - a_1(b_1 b_2 + z)(0))}{a_1(b_1 b_2 + z)(1) - ((1)(b_1 b_2 + z) + b_2 a_1)c'_1} \right| = \left| \frac{1589}{588} \right| = 2.$$

$$c'_3 = c_{3,2} = \frac{z(k_1(a_1)k_3(b_1)k_0(c'_2) - (k_0(a_2)k_3(b_1) + k_2(b_2)k_1(a_1))k_1(c'_1))}{(k_0(a_2)k_3(b_1) + k_2(b_2)k_1(a_1))k_2(c'_1) - k_1(a_1)k_3(b_1)k_1(c'_2)} = \frac{z(a_1(b_1 b_2 b_3 + z b_1 + z b_3)(1) - ((1)(b_1 b_2 b_3 + z b_1 + z b_3) + (b_2 b_3 + z) a_1) c'_1)}{(1)(b_1 b_2 b_3 + z b_1 + z b_3) + (b_2 b_3 + z) a_1 (c'_1 c'_2 + z) - a_1(b_1 b_2 b_3 + z b_1 + z b_3) c'_2} =$$

$$\frac{77}{31} \text{ (to be treated as C.F.)} = [2,3,2,2,19,2,4,5,2,3,8,4,4,2,2,9,5,7]_{7/4}$$

$$\text{Then } [2;2]_{7/4} + [2;9,5,7]_{7/4} = [4;1,2,2,3,2,2,19,2,4,5,2,3,8,4,4,2,2,9,5,7]_{7/4}$$

Example 3.3: Find $[4]_{7/4} + [0;7,5]_{7/4}$

Solution: Let $[4]_{7/4} = [a_0]_z$ and $[0;7,5]_{7/4} = [b_0; b_1, b_2]_z$, we get $m = 0, n = 2$ ($m < n$) and from definition (3.2b), we have $[4]_{7/4} + [0;7,5]_{7/4} = [a_0]_z + [b_0; b_1, b_2]_{7/4} = [c_0; b_1, b_2]_{7/4}$, where

$$c_0 = a_0 + b_0 = 4 + 0 = 4.$$

$$\text{Then } [4]_{7/4} + [0;7,5]_{7/4} = [4;7,5]_{7/4}$$

IV. SUBTRACTION OPERATION FOR TWO CONTINUED FRACTIONS

Definition 4.1: Let $[a_0; a_1, \dots, a_n]_z$ be a C.F then we can define additive inverses by $-[a_0; a_1, \dots, a_n]_z = [b'_0; b'_n]_z$, where

$$b'_0 = -1 - a_0 = f_0.$$

$$b'_n = \frac{z k_n(a_1)}{k_n(a_1) - z k_{n-1}(a_2)} \text{ (to be treated as C.F.)}$$

$$= [f_1; f_2, \dots, f_l]_z.$$

Therefore $-[a_0; a_1, \dots, a_n]_z = [b'_0; b'_n]_z = [f_0; f_1, \dots, f_l]_z$.

Example 4.1: Find $-[2;1,3]_{2/3}$.

Solution: Let $[2;1,3]_{2/3} = [a_0; a_1, a_2]_z$, ($n = 2$) from definition 4.1, we have: $[-2;1,3]_{2/3} = [b'_0; b'_2]_z$, where

$$b'_0 = -1 - a_0 = -1 - 2 = -3.$$

$$b'_2 = \frac{z k_2(a_1)}{k_2(a_1) - z k_1(a_2)} = \frac{22}{15} \text{ (to be treated as C.F.)}$$

$$= [1; 1, 1, 1, 3, 2]_{2/3}$$

$$\text{Then } -[2;1,3]_{2/3} = [-3; 1, 1, 1, 1, 3, 2]_{2/3}$$

Definition 4.2: If $[a_0; a_1, \dots, a_m]_z$ and $[b_0; b_1, \dots, b_n]_z$ are two C.F then we define subtraction $[a_0; a_1, \dots, a_m]_z - [b_0; b_1, \dots, b_n]_z$ by the addition $[a_0; a_1, \dots, a_m]_z + [f_0; f_1, \dots, f_l]_z$ where $[f_0; f_1, \dots, f_l]_z$ give by definition 4.1. That is

$$(1) \text{ If } m = l \text{ then } [a_0; a_1, \dots, a_m]_z + [f_0; f_1, \dots, f_m]_z = [c_0; c_1, \dots, c_m]_z \text{ (4.2a)}$$

where c_0, c_1, \dots, c_m as give in (3.2a)

(2) If $m \neq l, (l < m)$ then

$$[a_0; a_1, \dots, a_l, a_{l+1}, \dots, a_m]_z + [f_0; f_1, \dots, f_l]_z = [c'_0; c'_1, \dots, c'_l, c'_{l+1}, \dots, c'_m]_z \text{ (4.2b)}$$

where c'_0, c'_1, \dots, c'_m as give in (3.2b).

Example 4.2: Find $[1;1,2]_{5/3} - [0;1,2]_{5/3}$

Solution: (1) first we find $-[0;1,2]_{5/3}$. Let $[0;1,2]_{5/3} = [a_0; a_1, a_2]_{5/3}$, ($n = 2$). From definition 4.1, we have

$$-[0;1,2]_{5/3} = [b'_0; b'_2]_{5/3}, \text{ where}$$

$$b'_0 = -1 - a_0 = -1 - 0 = -1$$

$$b'_2 = \frac{z k_2(a_1)}{k_2(a_1) - z k_1(a_2)} = \frac{z(a_1 a_2 + z)}{(a_1 a_2 + z) - z(a_2)} = \frac{55}{3} = [18;5]_{5/3}.$$

$$\text{Then } -[0;1,2]_{5/3} = [-1; 18, 5]_{5/3}.$$

(2) To find $[1;1,2]_{5/3} + [-1;18,5]_{5/3}$, we let $[1;1,2]_{5/3} = [a_0; a_1, a_2]_z$ and $[-1,18,5]_{5/3} = [f_0; f_1, f_2]_z$, ($m = n = 2$) from (3.2a), we have

$$[1;1,2]_{5/3} + [-1;18,5]_{5/3} = [a_0; a_1, a_2]_z + [f_0; f_1, f_2]_z = [c_0; c_1, c_2]_z, \text{ where } c_2 \text{ is the last term and}$$

$$c_0 = a_0 + f_0 = 1 + (-1) = 0,$$

$$c_1 = \left| \frac{a_1 f_1}{a_1 + f_1} \right| = \left| \frac{1 \cdot 18}{1 + 18} \right| = \left| \frac{18}{19} \right| = 0,$$

$$c_2 = \frac{z(a_2(f_1 f_2 + z) + f_2(a_1 a_2 + z))}{(a_1 a_2 + z)(f_1 f_2 + z) - (a_2(f_1 f_2 + z) + f_2(a_1 a_2 + z))c_1} = \frac{3025}{3025} = 1,$$

Then $[1; 1, 2]_{5/3} + [-1; 18, 5]_{5/3} = [0; 0, 1]_{5/3} = 1$ (by lemma 2.1)

V. CONCLUSION

This paper is a sequel to our previous work in which we found the operations of the simple continued fractions. In the previous work, [8] we discovered the definitions of addition, subtractions, multiplication and division operations of the simple continued fractions. In this paper we defined the addition and subtractions of continued fractions with positive non Integer numerators. The study of the continued fractions with positive non Integer numerators is the first step to understand the continued fractions with polynomial numerators.

REFERENCES

- [1] Alan Baker (1986). "A concise introduction to theory of numbers". Cambridge University press,.
- [2] Burton, D. M. (2006). "Elementary number theory". Tata McGraw-Hill Education.
- [3] Greene, J., & Schmiege, J. (2017). "Continued Fractions with Non-Integer Numerators". J. Integer Seq., 20(1), 17-1.
- [4] Hardy, G. H., Wright, E. M., Heath-Brown, R., Silverman, J., & Wiles, A. (2008). "An Introduction to the Theory of Numbers". Oxford University Press.
- [5] LeVeque, W. J. (2014). "Fundamentals of Number Theory". Courier Corporation.
- [6] Mugassabi S. and Amsheri S. (2020) . "The Multiplication and Division of Simple Continued Fractions". IJISM International Journal of Innovation in Science and Mathematics 8 (3): (2020) 2347–9051.
- [7] Mugassabi, Souad I., and A. S. Elmabrok. (2019). "The Power of Simple Continued Fractions." EPH-International Journal of Applied Science (ISSN: 2208-2182) 1.1: 824-833.
- [8] Mugassabi S. and Mistiri F. (2015). "The Elementary Arithmetic Operators of Continued Fractions". Am-Euras. J. Sci.Res.,10 (5): 251 - 263.
- [9] C.D. Olds. (1963). "Continued Fractions". 5th Edition, New York: Random House, Inc.